

Section A

Q1 Find the 9th term from the end (towards the first term) of the A.P. 5, 9, 13, ..., 185.

Ans:- Common difference, d , of the AP = $9 - 5 = 4$

Last term, l , of the AP = 185

We know that the n^{th} term from the end of an AP is given by $l - (n - 1)d$.

Thus, the 9th term from the end is

$$185 - (9 - 1)4$$

$$= 185 - 4 \times 8$$

$$= 185 - 32 = 153$$

Q2 Cards marked with number 3, 4, 5, ..., 50 are placed in a box and mixed thoroughly. A card is drawn at random from the box. Find the probability that the selected card bears a perfect square number.

Ans:- It is given that the box contains cards marked with numbers 3, 4, 5, ..., 50.

\therefore Total number of outcomes = 48

Between the numbers 3 and 50, there are six perfect squares, i.e. 4, 9, 16, 25, 36 and 49.


\therefore Number of favourable outcomes = 6

\therefore Probability that a card drawn at random bears a perfect square

$$= \frac{\text{Number of favourable outcomes}}{\text{Total number of outcomes}} = \frac{6}{48} = \frac{1}{8}$$

Q3 From an external point P, tangents PA and PB are drawn to a circle with centre O. If $\angle PAB = 50^\circ$, then find $\angle AOB$.

Ans:-



It is given that PA and PB are tangents to the given circle.

$\therefore \angle PAO = 90^\circ$ (Radius is perpendicular to the tangent at the point of contact.)

Now, $\angle PAB = 50^\circ$ (Given)

$\therefore \angle OAB = \angle PAO - \angle PAB = 90^\circ - 50^\circ = 40^\circ$

In $\triangle OAB$,

$OB = OA$ (Radii of the circle)


$\therefore \angle OAB = \angle OBA = 40^\circ$ (Angles opposite to equal sides are equal.)

Now,

$\angle AOB + \angle OAB + \angle OBA = 180^\circ$ (Angle sum property)

$\Rightarrow \angle AOB = 180^\circ - 40^\circ - 40^\circ = 100^\circ$

Q4 In Fig. 1, AB is a 6 m high pole and CD is a ladder inclined at an angle of 60° to the horizontal and reaches up to a point D of pole. If $AD = 2.54$ m, find the length of the ladder. (use $\sqrt{3} = 1.73$)



Ans:- In the given figure,

$AB = AD + DB = 6$ m

Given: $AD = 2.54$ m

$$\Rightarrow 2.54 \text{ m} + DB = 6 \text{ m}$$

$$\Rightarrow DB = 3.46 \text{ m}$$

Now, in the right triangle BCD,

$$\frac{BD}{CD} = \sin 60^\circ$$

$$\Rightarrow \frac{3.46 \text{ m}}{CD} = \frac{\sqrt{3}}{2}$$

$$\Rightarrow \frac{3.46 \text{ m}}{CD} = \frac{1.73}{2}$$

$$\Rightarrow CD = \frac{2 \times 3.46 \text{ m}}{1.73}$$

$$\Rightarrow CD = 4 \text{ m}$$

Thus, the length of the ladder CD is 4 m.

Section B

Q5 Find the ratio in which y-axis divides the line segment joining the points A(5, -6) and B(-1, -4). Also find the coordinates of the point of division.

Ans:- Let $(0, \alpha)$ be a point on the y-axis dividing the line segment AB in the ratio $k : 1$.

Now, using the section formula, we get

$$(0, \alpha) = \left(\frac{-k+5}{k+1}, \frac{4k-6}{k+1} \right)$$

$$\Rightarrow \frac{-k+5}{k+1} = 0,$$

$$-4k - 6k + 1 = \alpha$$

Now,

$$\frac{-k+5}{k+1} = 0$$

$$\Rightarrow -k+5 = 0$$

$$\Rightarrow k = 5$$

Also,

$$\frac{4k-6}{k+1} = \alpha$$

$$\Rightarrow \frac{-4 \times 5 - 6}{5+1} = \alpha$$

$$\Rightarrow \alpha = \frac{-26}{6}$$

$$\Rightarrow \alpha = -\frac{13}{3}$$

Thus, the y-axis divides the line segment in the ratio $k : 1$, i.e. $5 : 1$.

Also, the coordinates of the point of division are $(0, a)$, i.e. $\left(0, -\frac{13}{3}\right)$.

Q6 If $x = \frac{2}{3}$ and $x = -3$ are roots of the quadratic equation $ax^2 + 7x + b = 0$, find the values of a and b .

Ans:- The given equation is $ax^2 + 7x + b = 0$. Its roots are given as -3 and $\frac{2}{3}$.

Now,
$$\text{Sum of the roots} = \frac{-(\text{Coefficient of } x)}{\text{Coefficient of } x^2}$$

$$\Rightarrow -3 + \frac{2}{3} = \frac{-(7)}{a}$$

$$\Rightarrow \frac{-9+2}{3} = \frac{-7}{a}$$

$$\Rightarrow \frac{-7}{3} = \frac{-7}{a}$$

$$\Rightarrow a = 3$$

Also, Product of the roots = $\frac{\text{Constant term}}{\text{Coefficient of } x^2}$

$$\Rightarrow -3 \times \frac{2}{3} = \frac{b}{a}$$

$$\Rightarrow -2 = \frac{b}{3}$$

$$\Rightarrow b = -6$$

Thus, the values of a and b are 3 and -6 , respectively

Q7 The x -coordinate of a point P is twice its y -coordinate. If P is equidistant from $Q(2, -5)$ and $R(-3, 6)$, find the coordinates of P .

Ans:- Let the y -coordinate of the point P be a .

Then, its x -coordinate will be $2a$.

Thus, the coordinates of the point P are $(2a, a)$.

It is given that the point $P(2a, a)$ is equidistant from $Q(2, -5)$ and $R(-3, 6)$.

Thus, we have $\sqrt{(2a-2)^2 + (a-(-5))^2} = \sqrt{(2a-(-3))^2 + (a-6)^2}$

$$\Rightarrow \sqrt{(2a-2)^2 + (a+5)^2} = \sqrt{(2a+3)^2 + (a-6)^2}$$

$$\Rightarrow \sqrt{4a^2 + 4 - 8a + a^2 + 25 + 10a} = \sqrt{4a^2 + 9 + 12a}$$

$$\Rightarrow \sqrt{5a^2 + 2a + 29} = \sqrt{5a^2 + 45}$$

$$5a^2 + 2a + 29 = 5a^2 + 45 \quad \text{Squaring both sides, we get}$$


$$\Rightarrow 5a^2 + 2a - 5a^2 = 45 - 29$$

$$\Rightarrow 2a = 16$$

$$\Rightarrow a = 8$$

Thus, the coordinates of the point P are $(16, 8)$, i.e. $(2 \times 8, 8)$.

Q8 In Fig. 2, a circle is inscribed in a $\triangle ABC$, such that it touches the sides AB , BC and CA at points D , E and F respectively. If the lengths of sides AB , BC and CA are 12 cm, 8 cm and 10 cm respectively, find the lengths of AD , BE and CF .



Ans:- It is given that

$$AB = 12 \text{ cm}$$

$$\Rightarrow AD + BD = 12 \text{ cm} \dots(1)$$

$$BC = 8 \text{ cm}$$

$$\Rightarrow BE + CE = 8 \text{ cm} \dots(2)$$

$$CA = 10 \text{ cm}$$

$$\Rightarrow AF + CF = 10 \text{ cm} \dots(3)$$

CF and CE act as tangents to the circle from the external point C.

It is known that the lengths of tangents drawn from an external point to a circle are equal.

$$\therefore CF = CE \dots(4)$$

Similarly, AF and AD act as tangents to the circle from the external point A.

$$\therefore AF = AD \dots(5)$$

Also, BD and BE act as tangents to the circle from the external point B.

$$\therefore BD = BE \dots(6)$$

Using (4) and (2), we get

$$BE + CF = 8 \text{ cm} \dots(7)$$

Using (5) and (3), we get

$$AD + CF = 10 \text{ cm} \dots(8)$$

Using (6) and (1), we get

$$AD + BE = 12 \text{ cm} \dots(9)$$

Adding (7), (8) and (9), we get

$$BE + CF + AD + CF + AD + BE = 8 \text{ cm} + 10 \text{ cm} + 12 \text{ cm}$$

$$\Rightarrow 2AD + 2BE + 2CF = 30 \text{ cm}$$

$$\Rightarrow 2(AD + BE + CF) = 30 \text{ cm}$$

$$\Rightarrow AD + BE + CF = 15 \text{ cm} \dots(10)$$

Subtracting (7) from (10), we get

$$AD + BE + CF - BE - CF = 15 \text{ cm} - 8 \text{ cm}$$

$$\Rightarrow AD = 7 \text{ cm}$$

Subtracting (8) from (10), we get

$$AD + BE + CF - AD - CF = 15 \text{ cm} - 10 \text{ cm}$$

$$\Rightarrow BE = 5 \text{ cm}$$


Subtracting (9) from (10), we get

$$AD + BE + CF - AD - BE = 15 \text{ cm} - 12 \text{ cm}$$

$$\Rightarrow CF = 3 \text{ cm}$$

Thus, the lengths of AD, BE and CF are 7 cm, 5 cm and 3 cm, respectively.

Q9 In Fig. 3, AP and BP are tangents to a circle with centre O, such that AP = 5 cm and $\angle APB = 60^\circ$. Find the length of chord AB.



Ans:- PA and PB are tangents drawn to the given circle from an external point P.

It is known that the lengths of the tangents drawn from an external point to a circle are equal.

$$\therefore PA = PB$$

In $\triangle PAB$, sides PA and PB are of the same length.

Hence, $\triangle PAB$ is isosceles, with $PA = PB$ and $\angle PAB = \angle PBA = x$ (say).

It is given that

$$\angle APB = 60^\circ$$

We know that the sum of the angles of a triangle is 180° .

In $\triangle PAB$,

$$\angle PAB + \angle PBA + \angle APB = 180^\circ$$

$$\therefore x + x + 60^\circ = 180^\circ$$

$$\Rightarrow 2x = 120^\circ$$

$$\Rightarrow x = 60^\circ$$

Thus,

$$\angle PAB = \angle PBA = \angle APB = 60^\circ$$

Since all angles of $\triangle PAB$ are of the same measure, $\triangle PAB$ is equilateral, with $AP = BP = AB$.

It is given that

$$AP = 5 \text{ cm}$$

$$\therefore AB = AP = 5 \text{ cm}$$

Thus, the length of the chord AB is 5 cm.

Q10 How many terms of the A.P. 65, 60, 55, be taken so that their sum is zero?

Ans:- The given AP is 65, 60, 55, ...

First term of the AP = 65

Common difference = 60 - 65 = -5

Let the sum of the first x terms of the AP be 0.

Sum of first x terms =

$$\Rightarrow \frac{x}{2} [130 + (-5x + 5)] = 0$$

$$\Rightarrow x(130 - 5x + 5) = 0$$

$$\Rightarrow x(135 - 5x) = 0$$

Now, either $x = 0$ or $135 - 5x = 0$.

Since the number of terms cannot be 0, $x \neq 0$.

$$\therefore 135 - 5x = 0$$

$$\Rightarrow 135 = 5x$$

$$\Rightarrow x = 27$$

Thus, the sum of the first 27 terms of the AP is 0.

Section C

Q11 A well of diameter 4 m is dug 21 m deep. The earth taken out of it has been spread evenly all around it in the shape of a circular ring of width 3 m to form an embankment. Find the height of the embankment.

Ans:- Let r and h be the radius and depth of the well, respectively.

$$\therefore r = \frac{4}{2} = 2 \text{ m and } h = 21 \text{ m}$$

Let R and H be the outer radius and height of the embankment, respectively.

$$\therefore R = r + 3 = 2 + 3 = 5 \text{ m}$$

Now,

Volume of the earth used to form the embankment = Volume of the earth dug out of the well

$$\pi(R^2 - r^2)H = \pi r^2 h$$

$$\Rightarrow H = \frac{r^2 h}{R^2 - r^2}$$

$$\Rightarrow H = \frac{2^2 \times 21}{5^2 - 2^2} = 4 \text{ m}$$

Thus, the height of the embankment is 4 m.

Q12 If the sum of first 7 terms of an A.P. is 49 and that of its first 17 terms is 289, find the sum of first n terms of the A.P.

Ans:- Let the first term and the common difference of the given AP be a and d , respectively.

Sum of the first 7 terms, $S_7 = 49$

We know

$$S = \frac{n}{2} [2a + (n-1)d]$$

$$\Rightarrow \frac{7}{2}(2a + 6d) = 49$$

$$\Rightarrow \frac{7}{2} \times 2(a + 3d) = 49$$

$$\Rightarrow a + 3d = 7 \quad \dots(1)$$

Sum of the first 17 terms, $S_{17} = 289$

$$\Rightarrow \frac{17}{2}(2a + 16d) = 289$$

$$\Rightarrow \frac{17}{2} \times 2(a + 8d) = 289$$

$$\Rightarrow a + 8d = \frac{289}{17} = 17$$

$$\Rightarrow a + 8d = 17 \quad \dots(2)$$

Subtracting (2) from (1), we get

$$5d = 10$$

$$\Rightarrow d = 2$$

Substituting the value of d in (1), we get

$$a = 1$$

Now,

Sum of the first n terms is given by

$$S_n = \frac{n}{2} [2a + (n-1)d]$$

$$= \frac{n}{2} [2 \times 1 + 2(n-1)]$$


$$= n(1 + n - 1) = n^2$$

Therefore, the sum of the first n terms of the AP is n^2 .

Q13 The angles of depression of the top and bottom of a 50 m high building from the top of a tower are 45° and 60° respectively. Find the height of the tower and the horizontal distance between the tower and the

building.

Ans: 60°



Let the height of the tower AB be h m and the horizontal distance between the tower and the building BC be x m.

So,

$AE = (h - 50)$ m

$$\tan 45^\circ = \frac{AE}{ED}$$

$$\Rightarrow 1 = \frac{h - 50}{x}$$

$$\Rightarrow x = h - 50 \quad \dots(1)$$

In $\triangle ABC$,

$$\tan 60^\circ = \frac{AB}{BC}$$

$$\Rightarrow \sqrt{3} = \frac{h}{x}$$

$$\Rightarrow x\sqrt{3} = h \quad \dots(2)$$

Using (1) and (2), we get

$$x = \sqrt{3}x - 50$$

$$\Rightarrow x(\sqrt{3} - 1) = 50$$

$$\Rightarrow x = \frac{50(\sqrt{3} + 1)}{2} = 25 \times 2.73 = 68.25 \text{ m}$$

Substituting the value of x in (1), we get

$$68.25 = h - 50$$


$$\Rightarrow h = 68.25 + 50$$

$$\Rightarrow h = 118.25 \text{ m}$$


Hence, the height of tower is 118.25 m and the horizontal distance between the tower and the building is 68.25 m.

Q14 In Fig. 4, ABCD is a square of side 14 cm. Semi-circles are drawn with each side of square as diameter. Find the area of the shaded region. (

$$\text{use } \pi = \frac{22}{7} \text{)}$$



Ans:-



Let the four shaded regions be I, II, III and IV and the centres of the semicircles

be P, Q, R and S, as shown in the figure.

It is given that the side of the square is 14 cm.

Now,

Area of region I + Area of region III = Area of the square – Areas of the semicircles with centres S and Q.

$$= 14 \times 14 - 2 \times \frac{1}{2} \times \pi \times 7^2 \quad (? \text{ Radius of the semicircle} = 7 \text{ cm})$$

$$= 196 - 49 \times \frac{22}{7}$$

$$= 196 - 154$$

$$= 42 \text{ cm}^2$$

Similarly,

Area of region II + Area of region IV = Area of the square – Areas of the semicircles with centres P and R.

$$= 14 \times 14 - 2 \times \frac{1}{2} \times \pi \times 7^2 \quad (? \text{ Radius of the semicircle} = 7 \text{ cm})$$

$$= 196 - 49 \times \frac{22}{7}$$

$$= 196 - 154$$

$$= 42 \text{ cm}^2$$

Thus,


Area of the shaded region = Area of region I + Area of region III + Area of region II + Area of region IV

$$= 42 \text{ cm}^2 + 42 \text{ cm}^2$$


$$= 84 \text{ cm}^2$$

Q15 In Fig. 5, is a decorative block, made up two solids – a cube and a hemisphere. The base of the block is a cube of side 6 cm and the hemisphere fixed on the top has diameter of 3.5 cm. Find the total surface

area of the block. (use $\pi = \frac{22}{7}$)



Ans:-



Surface area of the block = Total surface area of the cube - Base area of the hemisphere + Curved surface area of the hemisphere

$$= 6 \times (\text{Edge})^2 - \pi r^2 + 2\pi r^2$$


$$= (6^2 + \pi r^2)$$

$$= \left(216 + \frac{22}{7} \times \frac{3.5}{2} \times \frac{3.5}{2} \right)$$


$$= (216 + 9.625)$$

$$= 225.625 \text{ cm}^2$$

Q16 In Fig. 6, ABC is a triangle coordinates of whose vertex A are $(0, -1)$. D and E respectively are the mid-points of the sides AB and AC and their coordinates are $(1, 0)$ and $(0, 1)$ respectively. If F is the mid-point of BC, find the areas of



Ans:-



(x_2, y_2) and (x_3, y_3) ,

Let the coordinates of B and C be (x_2, y_2) and (x_3, y_3) , respectively.
D is the midpoint of AB.

$$\text{So, } (1, 0) = \left(\frac{x_2 + 0}{2}, \frac{y_2 - 1}{2} \right)$$

$$\Rightarrow 1 = \frac{x_2}{2} \text{ and } 0 = \frac{y_2 - 1}{2}$$

$$\Rightarrow x_2 = 2 \text{ and } y_2 = 1$$

Thus, the coordinates of B are $(2, 1)$.

Similarly, E is the midpoint of AC.

So,

$$(0, 1) = \left(\frac{x_3 + 0}{2}, \frac{y_3 - 1}{2} \right)$$

$$\Rightarrow 0 = \frac{x_3}{2} \text{ and } 1 = \frac{y_3 - 1}{2}$$

$$\Rightarrow x^3 = 0 \text{ and } y^3 = 3$$

Thus, the coordinates of C are (0, 3).

Also, F is the midpoint of BC. So, its coordinates are

$$\left(\frac{2+0}{2}, \frac{1+3}{2} \right) = (1, 2)$$

Now,

Area of a triangle

$$= \frac{1}{2} [x_1(y_2 - y_3) + x_2(y_3 - y_1) + x_3(y_1 - y_2)]$$

Thus, the area of $\triangle ABC$ is

$$\frac{1}{2} [0(1-3) + 2(3+1) + 0(-1-1)]$$

$$= \frac{1}{2} \times 8$$

= 4 square units

And the area of $\triangle DEF$ is


$$\frac{1}{2} [1(1-2) + 0(2-0) + 1(0-1)]$$

$$= \frac{1}{2} \times (-2)$$

= 1 square unit (Taking the numerical value, as the area cannot be negative)

Q17 In Fig. 7, are shown two arcs PAQ and PBQ. Arc PAQ is a part of circle with centre O and radius OP while arc PBQ is a semi-circle drawn on PQ as diameter with centre M. If $OP = PQ = 10$ cm show that area of shaded region

is $25 \left(\sqrt{3} - \frac{\pi}{6} \right) \text{ cm}^2$




Ans:- Given: $OP = OQ = 10$ cm

It is known that tangents drawn from an external point to a circle are equal in length.

So,

$OP = OQ = 10$ cm



Therefore, $\triangle POQ$ is an equilateral triangle.

$\Rightarrow \angle POQ = 60^\circ$

Now,

Area of part II = Area of the sector - Area of the equilateral triangle POQ

$$= \frac{\angle POQ}{360^\circ} \times \pi r^2 - \frac{\sqrt{3}}{4} \times (10)^2$$

$$= \frac{60^\circ}{360^\circ} \times \pi (10^2) - \frac{\sqrt{3}}{4} \times (10)^2$$

$$= 100 \left(\frac{\pi}{6} - \frac{\sqrt{3}}{4} \right) \text{ sq units}$$

Area of the semicircle on diameter PQ = Area of part II + Area of part III

$$= \frac{1}{2} \times \pi (5^2) = \frac{25}{2} \pi \text{ sq units}$$

\therefore Area of the shaded region (part III)

$$= \frac{25}{2} \pi - \frac{100}{6} \left(\frac{\pi}{6} - \frac{\sqrt{3}}{4} \right)$$

$$= \frac{25}{2} \pi - \frac{100}{6} \pi + 25\sqrt{3}$$

$$= 25\sqrt{3} - \frac{25}{6}\pi$$

$$= 25\left(\sqrt{3} - \frac{\pi}{6}\right) \text{sq units}$$

Hence proved.

Q18 A box consists of 100 shirts of which 88 are good, 8 have minor defects and 4 have major defects. Ramesh, a shopkeeper will buy only those shirts which are good but 'Kewal' another shopkeeper will not buy shirts with major defects. A shirt is taken out of the box at random. What is the probability that

(i) Ramesh will buy the selected shirt?

(ii) 'Kewal' will buy the selected shirt?

Ans:- Total number of shirts = 100

Number of good shirts = 88

Number of shirts with minor defects = 8

Number of shirts with major defects = 4

(i) It is given that Ramesh will buy only good shirts.

$$\therefore P(\text{Selected shirt is good shirt}) = \frac{88}{100} = \frac{22}{25}$$

$$\Rightarrow \text{Probability that Ramesh will buy the selected shirt} = \frac{22}{25}$$

(ii) It is given that Kewal will not buy shirts with major defects. So, Kewal will buy good shirts and shirts with minor defects.

Total number of good shirts and shirts with minor defects = $88 + 8 = 96$

$$\therefore P(\text{Selected shirt is good or shirt with minor defect}) = \frac{96}{100} = \frac{24}{25}$$

$$\Rightarrow \text{Probability that Kewal will buy the selected shirt} = \frac{24}{25}$$

Q19 Solve the following quadratic equation for x:

$$x^2 + \left(\frac{a}{a+b} + \frac{a+b}{a}\right)x + 1 = 0$$

Ans:- Given:

$$x^2 + \left(\frac{a}{a+b} + \frac{a+b}{a}\right)x + 1 = 0$$

let

$$\frac{a}{a+b} be$$

Thus, the equation becomes

$$x^2 + \left(t + \frac{1}{t}\right)x + 1 = 0.$$

$$\Rightarrow x^2 + \left(t + \frac{1}{t}\right)x + 1 = 0$$

$$\Rightarrow x^2 + \left(\frac{t^2 + 1}{t}\right)x + 1 = 0$$

$$\Rightarrow tx^2 + (t^2 + 1)x + t = 0$$

$$\Rightarrow tx^2 + t^2x + x + t = 0$$

$$\Rightarrow (tx^2 + t^2x) + (x + t) = 0$$

$$\Rightarrow tx(x + t) + 1(x + t) = 0$$

$$\Rightarrow (tx + 1)(x + t) = 0$$

$$\Rightarrow tx + 1 = 0, x + t = 0$$

$$\Rightarrow x = -\frac{1}{t}, x = -t$$


$$\Rightarrow x = -\frac{1}{\frac{a}{a+b}}, x = -\frac{a}{a+b}$$

$$\Rightarrow x = -\frac{a+b}{a}, x = -\frac{a}{a+b}$$

Q20 A toy is in the form of a cone of base radius 3.5 cm mounted on a hemisphere of base diameter 7 cm. If the total height of the toy is 15.5 cm,

find the total surface area of the toy. *use $\pi = \frac{22}{7}$*

Ans:-



Let r and h be the radius and height of the cone mounted on the hemisphere, respectively.

Suppose R be the radius of the hemisphere.

Now,

$$r = R = 3.5 \text{ cm}$$

Height of the cone + Radius of the hemisphere = Total height of the toy

$$\therefore h + 3.5 \text{ cm} = 15.5 \text{ cm}$$

$$\Rightarrow h = 15.5 - 3.5 = 12 \text{ cm}$$

Let l be the slant height of the cone.

$$\therefore l^2 = r^2 + h^2$$

$$\Rightarrow l^2 = \left(\frac{7}{2}\right)^2 + (12)^2$$

$$= \frac{49}{4} + 144 = \frac{625}{4}$$

$$\Rightarrow l = \frac{25}{2} \text{ cm}$$

Total surface area of the toy

= Curved surface area of the cone + Curved surface area of the hemisphere

$$= \pi r l + 2\pi r^2$$

$$= \pi r (l + 2r)$$

$$= \frac{22}{7} \times \frac{7}{2} \times \left(\frac{25}{2} + 2 \times \frac{7}{2}\right)$$

$$= \frac{22}{7} \times \frac{7}{2} \times \frac{39}{2} = 214.5 \text{ cm}^2$$

Section D

Q21 A passenger, while boarding the plane, slipped from the stairs and got hurt. The pilot took the passenger in the emergency clinic at the airport for treatment. Due to this, the plane got delayed by half an hour. To reach the destination 1500 km away in time, so that the passengers could catch the connecting flight, the speed of the plane was increased by 250 km/hour than the usual speed. Find the usual speed of the plane.

What value is depicted in this question?

Ans:- Let the usual speed of the plane be x km/h.

Let the time taken by the plane to reach the destination be t_1 .

$$\therefore t_1 = \frac{1500}{x}$$

To reach the destination on time, the speed of the plane was increased to $(x + 250)$ km/h.

$$\therefore t_2 = \frac{1500}{x+250}$$

Given: $t_1 - t_2 = 30 \text{ min}$

Now,

$$\frac{1500}{x} - \frac{1500}{x+250} = \frac{30}{60}$$

$$\Rightarrow \frac{1500(x+250-x)}{x(x+250)} = \frac{1}{2}$$

$$\Rightarrow 750000 = x^2 + 250x$$

$$\Rightarrow x^2 + 250x - 750000 = 0$$

$$\Rightarrow x^2 + 250x - 750000 = 0$$

On solving the equation, we get

$$x = 750$$


Thus,

Usual speed of the plane = 750 km/h

The value depicted in this question is that of humanity. The pilot has set an example of a good and responsible citizen of the society.

Q22 In Fig. 8, O is the centre of a circle of radius 5 cm. T is a point such that OT = 13 cm and OT intersects circle at E. If AB is a tangent to the circle at E,

find the length of AB, where TP and TQ are two tangents to the circle.



Ans:- From the given figure, we have

$TP = TQ$ (Two tangents, drawn from an external point to a circle, have equal length.)

and

$\angle TQO = \angle TPO = 90^\circ$ (Tangent to a circle is perpendicular to the radius through the point of contact.)

In $\triangle TOQ$,

$$QT^2 + OQ^2 = OT^2$$

$$\Rightarrow QT^2 = 13^2 - 5^2 = 144$$

$$\Rightarrow QT = 12 \text{ cm}$$

Now,

$$OT - OE = ET = 13 - 5 = 8 \text{ cm}$$

Let $QB = x \text{ cm}$.

$\therefore QB = EB = x$ (Two tangents, drawn from an external point to a circle, have equal length.)

Also, $\angle OEB = 90^\circ$ (Tangent to a circle is perpendicular to the radius through the point of contact.)

In $\triangle TEB$,

$$EB^2 + ET^2 = TB^2$$

$$\Rightarrow x^2 + 8^2 = (12 - x)^2$$

$$\Rightarrow x^2 + 64 = 144 + x^2 - 24x$$

$$\Rightarrow 24x = 80$$


$$\Rightarrow x = \frac{80}{24} = \frac{10}{3}$$

$$\therefore AB = 2x = \frac{20}{3} \text{ cm}$$

Thus, the length of AB is $\frac{20}{3} \text{ cm}$.

Q23 A bird is sitting on the top of a 80 m high tree. From a point on the ground, the angle of elevation of the bird is 45° . The bird flies away horizontally in such a way that it remained at a constant height from the ground. After 2 seconds, the angle of elevation of the bird from the same point is 30° . Find the speed of flying of the bird. (Take $\sqrt{3} = 1.732$).

Ans:- Let P and Q be the two positions of the bird, and let A be the point of observation. Let ABC be the horizontal line through A.
Given: The angles of elevations of the bird in two positions P and Q from point A are 45° and 30° , respectively.



$$\therefore \angle PAB = 45^\circ \text{ and } \angle QAB = 30^\circ$$

Also,

$$PB = 80 \text{ m}$$

In $\triangle ABP$, we have

$$\tan 45^\circ = \frac{BP}{AB}$$

$$\Rightarrow 1 = \frac{80}{AB}$$

$$\Rightarrow AB = 80 \text{ m}$$

In $\triangle ACQ$, we have

$$\tan 30^\circ = \frac{CQ}{AC}$$

$$\Rightarrow \frac{1}{\sqrt{3}} = \frac{80}{AC}$$

$$\Rightarrow AC = 80\sqrt{3} \text{ m}$$

$$\therefore PQ = BC = AC - AB$$


$$= 80\sqrt{3} - 80 = 80(\sqrt{3} - 1) \text{ m}$$

So, the bird covers $80(\sqrt{3} - 1)$ m in 2 s.

Thus, speed of the bird is given by

$$\begin{aligned}
 &= \frac{80(\sqrt{3}-1)}{2} \text{ m/s} \\
 &= \frac{40(\sqrt{3}-1) \times 60 \times 60}{1000} \text{ km/h} \\
 &= 144(1.732-1) \text{ km/h} \\
 &= 105.408 \text{ km/h.}
 \end{aligned}$$

Q24 A game of chance consists of spinning an arrow on a circular board, divided into 8 equal parts, which comes to rest pointing at one of the numbers 1, 2, 3, ..., 8 (Fig. 9), which are equally likely outcomes. What is the probability that the arrow will point at (i) an odd number (ii) a number greater than 3 (iii) a number less than 9.



Ans:- Arrow can come to rest at any of the numbers 1, 2, 3, 4, 5, 6, 7 and 8.

Total number of events = 8

(i) There are four odd numbers 1, 3, 5 and 7.

Probability that the arrow will point at an odd number is given by

$$P(\text{Arrow point at odd number}) = \frac{4}{8} = \frac{1}{2}$$

(ii) There are five numbers greater than 3, that is, 4, 5, 6, 7 and 8.

Probability that the arrow will point at a number greater than 3 is given by

$$P(\text{Arrow point at a number greater than 3}) = \frac{5}{8}$$

(iii) All the numbers are less than 9.

Probability that the arrow will point at a number less than 9 is given by

$$P(\text{Arrow point at a number less than 9}) = \frac{8}{8} = 1$$

Q25 Prove that the area of a triangle with vertices $(t, t - 2)$, $(t + 2, t + 2)$ and $(t + 3, t)$ is independent of t .

Ans:- Let A $(t, t - 2)$, B $(t + 2, t + 2)$ and C $(t + 3, t)$ be the vertices of the given triangle.

We know that the area of the triangle having vertices

(x_1, y_1) , (x_2, y_2) and (x_3, y_3) is

$$\frac{1}{2} |x_1(y_2 - y_3) + x_2(y_3 - y_1) + x_3(y_1 - y_2)|.$$

\therefore Area of $\triangle ABC$

$$= \frac{1}{2} [x_1(y_2 - y_3) + x_2(y_3 - y_1) + x_3(y_1 - y_2)]$$

$$= \frac{1}{2} [t(t + 2 - t) + (t + 2)(t - t + 2) + (t + 3)(t - 2 - t - 2)]$$

$$= \frac{1}{2} (2t + 2t + 4 - 4t - 12)$$


$$= |-4|$$

$$= 4 \text{ square units}$$

Hence, the area of the triangle with given vertices is independent of t .

Q26 A bucket open at the top is in the form of a frustum of a cone with a capacity of 12308.8 cm^3 . The radii of the top and bottom circular ends are 20 cm and 12 cm, respectively. Find the height of the bucket and the area of metal sheet used in making the bucket. (use $\pi = 3.14$)

Ans:- Consider the following figure:



Given: Volume of the frustum is 12308.8 cm^3 .

Radii of the top and bottom are $r_1 = 20 \text{ cm}$ and $r_2 = 12 \text{ cm}$, respectively.
Volume of the frustum is given by

$$V = \frac{1}{3} \pi h (r_1^2 + r_2^2 + r_1 r_2)$$

$$12308.8 \times 3 = \pi h (20^2 + 12^2 + 20 \times 12)$$

$$12308.8 \times 3 = \pi h (400 + 144 + 240) \quad 12308.8 \times 3 = \pi h (784)$$

$$\frac{12308.8 \times 3}{3.14 \times 784} = h$$

$$\frac{3920 \times 3}{784} = h$$

$$15 \text{ cm} = h$$

Hence, height of the frustum is 15 cm.

Now,

Metal sheet required to make the frustum = Curved surface area + Area of the base of the frustum

Curved surface area of the frustum

$$= \pi (r_1 + r_2) l, \quad \text{where } l = \sqrt{h^2 + (r_1 - r_2)^2}$$

$$= \sqrt{225 + 64} = \sqrt{289} = 17 \text{ cm}$$

Curved surface area of the frustum

$$= \pi (20 + 12) 17$$


$$= 544 \times 3.14$$

$$= 1708.16 \text{ cm}^2$$


$$\text{Area of the base} = \pi 12^2 = 144 \times 3.14 = 452.16 \text{ cm}^2$$

$$\therefore \text{Metal sheet required to make the frustum} = 1708.16 + 452.16 = 2160.32 \text{ cm}^2$$

Q27 An elastic belt is placed around the rim of a pulley of radius 5 cm. (Fig. 10) From one point C on the belt, the elastic belt is pulled directly away from the centre O of the pulley until it is at P, 10 cm from the point O. Find the length of the belt that is still in contact with the pulley. Also find the shaded area. (use $\pi = 3.14$ and $\sqrt{3} = 1.73$)



Ans:-



Given: $OA = 5 \text{ cm}$

$OP = 10 \text{ cm}$

We know that the tangent at any point of a circle is perpendicular to the radius through the point of contact.

Therefore, $\triangle OAP$ is a right-angled triangle.

$$\Rightarrow \angle OAP = 90^\circ$$

Now,

$$OP^2 = OA^2 + AP^2$$

$$\Rightarrow 10^2 = 5^2 + AP^2$$

$$\Rightarrow AP^2 = 75$$

$$\Rightarrow AP = 5\sqrt{3} \text{ cm}$$

Also,

$$\cos \theta = \frac{OA}{OP} = \frac{5}{10}$$

$$\Rightarrow \cos \theta = \frac{1}{2}$$

$$\Rightarrow \theta = 60^\circ$$

Now, $\angle AOP = \angle BOP = 60^\circ$ ($\because \triangle OAP \cong \triangle OBP$)

$$\Rightarrow \angle AOB = 120^\circ$$

Length of the belt still in contact with the pulley = Circumference of the circle
 - Length of the arc ACB

$$= 2 \times 3.14 \times 5 - \frac{120^\circ}{360^\circ} \times 2 \times 3.14 \times 5$$

$$= 2 \times 3.14 \times 5 \times \left(1 - \frac{1}{3}\right)$$

$$= 2 \times 3.14 \times 5 \times \frac{2}{3}$$

20.93 cm (Approx.)

Now,

Area of $\triangle OAP$

$$= \frac{1}{2} \times AP \times OA = \frac{1}{2} \times 5\sqrt{3} \times 5$$

$$= \frac{25\sqrt{3}}{2} \text{ cm}^2$$

Similarly,

$$\text{Area of } \triangle OBP = \frac{25\sqrt{3}}{2} \text{ cm}^2$$

\therefore Area of $\triangle OAP$ + Area of $\triangle OBP$

$$= 25\sqrt{3} \text{ cm}^2 = 25 \times 1.73$$

$$= 43.25 \text{ cm}^2$$

$$\text{Area of sector OACB} = \frac{120^\circ}{360^\circ} \times 3.14 \times (5)^2$$

$$= \frac{1}{3} \times 3.14 \times 25 = 26.17 \text{ cm}^2 \text{ (Approx.)}$$

\therefore Area of the shaded region = (Area of $\triangle OAP$ + Area of $\triangle OBP$) - Area of the sector OACB

$$= 43.25 \text{ cm}^2 - 26.17 \text{ cm}^2$$

$$= 17.08 \text{ cm}^2 \text{ (Approx.)}$$

Q28 The sum of three numbers in A.P. is 12 and sum of their cubes is 288.

Find the numbers.

Ans:- Let the three numbers be $a - d$, a and $a + d$.

$$S_3 = 12$$

$$a - d + a + a + d = 12$$

$$\Rightarrow 3a = 12$$

$$\Rightarrow a = 4 \dots (1)$$

$$\begin{aligned}
 (a - d)^3 + a^3 + (a + d)^3 &= 288 \\
 \Rightarrow (4 - d)^3 + 4^3 + (4 + d)^3 &= 288 \\
 \Rightarrow 64 + d^3 + 48d + 12d^2 + 64 + 64 - d^3 - 48d + 12d^2 &= 288 \\
 \Rightarrow 24d^2 &= 96 \\
 \Rightarrow d &= \pm 2
 \end{aligned}$$

If $d = 2$, then the numbers are $(4 - 2)$, 4 and $(4 + 2)$, that is, 2 , 4 and 6 .

If $d = -2$, then the numbers are $(4 + 2)$, 4 and $(4 - 2)$, that is, 6 , 4 and 2 .


Q29 Prove that the lengths of tangents drawn from an external point to a circle are equal.

Ans:- Given: A circle with centre O , a point P lying outside the circle and PQ and PR as the two tangents

To prove: $PQ = PR$

Construction: Join OP , OQ and OR .

Proof:



In $\triangle OQP$ and $\triangle ORP$,

$\angle OQP = \angle ORP = 90^\circ$ (Tangent at any point of a circle is perpendicular to the radius through the point of contact.)

$OQ = OR$ (Radii)

$OP = OP$ (Common)

$\therefore \triangle OQP \cong \triangle ORP$ (By RHS congruency criterion)

$\Rightarrow PQ = PR$ (Corresponding parts of congruent triangles)

Q30 The time taken by a person to cover 150 km was $2\frac{1}{2}$ hours more than the time taken in the return journey. If he returned at a speed of 10 km/hour more than the speed while going, find the speed per hour in each direction.

Ans:- Let the speed of the person while going be x km/h.

\Rightarrow Speed of the person while returning = $(x + 10)$ km/h

Time taken by the person while going = Time taken by the person while

$$\begin{aligned} \therefore \frac{150 \text{ km}}{x \text{ km/h}} &= \frac{150 \text{ km}}{(x+10) \text{ km/h}} + \frac{5}{2} \\ \Rightarrow \frac{150}{x} - \frac{150}{x+10} &= \frac{5}{2} \\ \Rightarrow \frac{150x+1500-150x}{x(x+10)} &= \frac{5}{2} \\ \Rightarrow x^2 + 10x &= 1500 \times \frac{2}{5} = 600 \\ \Rightarrow x^2 + 10x - 600 &= 0 \\ \Rightarrow x^2 + 30x - 20x - 600 &= 0 \\ \Rightarrow x(x+30) - 20(x+30) &= 0 \end{aligned}$$

$$= (x+30)(x-20)=0$$

$$=x-20=0 \text{ or } x=30$$

$$=x=20, -30$$

Since speed cannot be negative, $x = 20$.

\Rightarrow Thus, speed of the person while going = 20 km/h

\Rightarrow Speed of the person while returning = $20 + 10 = 30$ km/h

Q31 Draw a triangle ABC with $BC = 7$ cm, $\angle B = 45^\circ$ and $\angle A = 105^\circ$. Then construct a triangle whose sides are times the corresponding sides of ΔABC .

Ans:- Given:

$$\angle B = 45^\circ$$

$$\angle A = 105^\circ$$

Sum of all interior angles in a triangle is 180° .

$$\angle A + \angle B + \angle C = 180^\circ$$

$$105^\circ + 45^\circ + \angle C = 180^\circ$$

$$\angle C = 180^\circ - 150^\circ$$

$$\angle C = 30^\circ$$

Following steps are involved in the construction of the required triangle:

Step 1

Draw a $\triangle ABC$ with side $BC = 7$ cm, $\angle B = 45^\circ$ and $\angle C = 30^\circ$.

Step 2

Draw a ray BX making an acute angle with BC on the opposite side of vertex A .

Step 3


Locate 5 points (as 5 is greater in 5 and 4) B_1, B_2, B_3, B_4 and B_5 on BX such that $BB_1, B_1B_2, B_2B_3, B_3B_4$ and B_4B_5 .

Step 4

Join B_5C . Draw a line through B_4 parallel to B_5C intersecting extended BC at C' .

Step 5

Through C' , draw a line parallel to AC intersecting BA at A' . $\triangle A'BC'$ is the required triangle.



Justification

The construction can be justified by proving that

$$A'B = \frac{4}{5} AB, BC' = \frac{4}{5} BC \text{ and } A'C' = \frac{4}{5} AC$$

In $\triangle ABC$ and $\triangle A'BC'$,

$\angle ABC = \angle A'BC'$ (Common)

$\angle ACB = \angle A'C'B$ (Corresponding angles)

$\therefore \triangle ABC \sim \triangle A'BC'$ (AA similarity criterion)

$$\frac{AB}{A'B} = \frac{BC}{BC'} = \frac{AC}{A'C'} \dots\dots (1)$$

In $\triangle BB_5C$ and $\triangle BB_4C'$,

$\angle B_4BC' = \angle B_5BC$ (Common)

$\angle BB_4C' = \angle BB_5C'$ (Corresponding angles)

$\therefore \triangle BB_4C' \sim \triangle BB_5C$ (AA similarity criterion)

$$\Rightarrow \frac{BC'}{BC} = \frac{4}{5} \dots\dots (2)$$

Comparing (1) and (2), we obtain

$$\frac{A'B}{AB} = \frac{BC'}{BC} = \frac{A'C'}{AC} = \frac{4}{5}$$

This justifies the construction.

Klass Studio